CONTRIBUTION OF GLACIER MELT WATER TO THE FLOW OF THE BOW RIVER

PHASE I Part D

Compilation of data and analysis of trends to the present

Final report

by

G J Young,
Wilfrid Laurier University

to

Government of the Province of Alberta
Department of Environmental Protection

June 1995

CONTRIBUTION OF GLACIER MELT WATER TO THE FLOW OF THE BOW RIVER

PHASE I Part D

Compilation of data and analysis of trends to the present

Final report

by

G J Young,
Wilfrid Laurier University

to

Government of the Province of Alberta
Department of Environmental Protection

June 1995

103-01848/

Table of Contents

List of figures and tables		**ii**
Acknowledgements		**iii**
Executive summary		**1**
Chapter 1	**Database I: Maps of the Bow River Basin above Banff and information contained in the Geographical Information System**	**3**
Chapter 2	**Database II: Streamflow records**	**12**
Chapter 3	**Database III: Meteorological records, snowcourse records and records of glacier mass balance**	**21**
Chapter 4	**Database IV: Glacier inventory and selected coverage by aerial photography**	**27**
Bibliography		**34**
Appendix	**Specific yield from sub-basins June, July, August, September, 1973-76**	**44**

List of figures and tables

1.1	The Bow River Basin above the Banff stream gauge (BB001), covering 2227 km^2, with sub-basins delineated.	4
1.2	The Bow River Basin above the Banff showing major drainage network and gauge sites for sub-basins.	5
1.3	Relief within the Bow River Basin above the Banff. 500 ft.	6
1.4	Forest distribution within the Bow River Basin above the Banff.	7
1.5	Glacier distribution within the Bow River Basin above the Banff.	8
1.6	Schematic of relative positions of stream gauges within the Bow River Basin above Banff	10
1.7	Hypsographic curves for the basin of the Bow River above Banff and for the Bow River above Lake Louise	11
2.1	Time line of availability of major series of discharge within the Bow River basin above Banff and in the adjacent Mistaya Basin	13
2.2	Mean annual flows of the Bow river at Banff 1910-1992	14
2.3	Mean monthly discharge of the Bow River at Banff 1909-1992	15
2.4	Mean monthly discharge of the Bow River at Banff and for the Bow River at Lake Louise 1965-1992	16
2.5	Discharge of the Bow River at Banff by month 1910-1992	17
2.6	Mean monthly flows of the Bow River at Banff 1973-76 as a percentage of the mean monthly flows 1911-1992	18
2.7	Bow River at Banff daily discharge 1973-1976	19
2.8	Standardized mean monthly discharges 1973-1976 for the Bow River at Banff, the Bow River below Hector Lake and Forty Mile Creek.	20
3.1	Annual temperature and precipitation records at Banff, 1895-1992	22
3.2	(a) Time line of availability of data and (b) location of selected snowcourses	23
3.3	April 1st water equivalents (mm) for the four snowcourses with the longest records	25
3.4	Peyto Glacier mass balance 1966-84, with extrapolations back to 1950	26
4.1	Glacier inventory base map of the Upper Bow River and adjacent basins from the Glacier ' tlas of Canada	28
4.2	Map of the Glacier Inventory of the Waputik Mountains from Müller et al 1967	29

Table 1.1	Characteristics of Bow River Basin above Banff and adjacent basins for comparative studies	9
Table 3.1	April 1st water equivalents (mm) for selected snowcourses 1937-1989	24
Table 4.1	Summary data from glacier inventory of the Waputik Mountains from Müller et al 1967	30-31
Table 4.2(a)	Air photographs from 1951-52	32
Table 4.2(b)	Air photographs from 1977 July / August	33
Table 4.2(c)	Air photographs from 1993 September	33

Acknowledgements

I am particularly indebted to Corinne Schuster, Rob South and Al Wallace for all their efforts in support of this work during the past year. Without their dedication to the project the compilation and illustration of the data sources would not have been possible.

Gordon J Young

Executive Summary

Within the Bow River Basin, Alberta, the demand for water is rising. Already, especially at critical times of the year (late summer) and in years of low flows, demands exceed Alberta's entitlements to water.

Very important questions to be answered at a time when demand is outstripping supply are:

- to what extent is the supply variable over years and decades;
- whether there are long term trends in climate leading to changing stream flows;
- to what extent supplies can be relied upon in the future.

The water supply is derived primarily from the mountain regions in the upper catchment areas where precipitation is relatively high. Much of the mountain precipitation is stored seasonally as snowpacks and over periods of decades and centuries in the form of glaciers. Glaciers provide meltwater primarily in late summer and, in years with relatively low precipitation, glaciers compensate for low flows through additional melting. ***Thus glaciers play a crucial role in water supply for it is precisely when demand is highest that they give their most valuable contributions.***

Over the last few decades glaciers have been in recession, releasing water into the rivers in addition to the annual water budget. This process may well continue and for a number of years there may be a very welcome augmentation of flows. However, as glacier tongues recede to ever higher (and colder) elevations, the contributions of glacier melt to streamflow must, of necessity diminish. ***It is crucial to the assessment of the available future water resources to understand the quantities and timing of release of glacier ice from long term storage and the rate at which that component of the water supply is likely to change in magnitude.***

It is important to note that the significance of the diminution of glacier extent, with concomitant reduction in water supply, might be completely overshadowed by long term trends in climate, resulting in changes in precipitation quantities or timing of melt periods, or by shorter term but highly significant variations in climate such as a series of years influenced by El Niño events.

While there is a great deal of uncertainty regarding climatic variation over both short and long term time scales, it is possible (at least approximately) to quantify the volume of ice being released from storage over periods of decades and thus to identify the significance of this component of streamflow.

It is the purpose of this research to identify the contribution of glacier ice melt to streamflow, and the changes in that component of the water balance through time in the past and into the future. The geographic extent of the analyses is confined to the catchment of the Bow River above Banff, a catchment within which there has been minimal human influence.

To address the problem the research has been broken down into three phases, the first of which, "Compilation of data and analysis of trends to the present", is the subject of this Report.

Several types of data have been compiled, inspected for quality and consistency and are presented in summary, primarily in graphical form. The data bases include:

- A series of maps using a Geographical Information System (Terrasoft) to delineate basin and sub-basin boundaries; streams and lakes; elevation; forest cover; glacier extent; and bare rock areas. By overlaying basin boundaries, terrain types and elevation, hypsographic curves may be derived; these will be an integral component in the further analysis and interpretation of the streamflow records.

- The streamflow records of the Bow River above Banff and some records of nearby catchments which will be used in subsequent analyses.

- Selected meteorological records (not including those of questionable accuracy and consistency), snowcourse records and records of glacier mass balance.

Analyses to be undertaken in further phases of the research include:

- Utilizing analyses already undertaken in the adjacent Mistaya Basin, transpose changes in glacier extent since 1966 and transpose the discharge records to the highly glacierized sections of the upper Bow River Basin.

- From maps, aerial photography and (perhaps) from satellite imagery, reconstruct the extent of glaciers in the Bow Basin in the time period 1966 to present and calculate volumetric change.

- Calculate the effects of volumetric change of glaciers on the hydrographs for the whole Bow Basin above Banff and on its sub-catchments.

- Construct future scenarios with smaller glaciers and with different combinations of climatic inputs in order to understand likely water supply conditions in the future.

CHAPTER 1 Database I: Maps of the Bow River Basin above Banff and information contained in the Geographical Information System

This chapter describes the layout of the Bow River Basin above Banff. The basin covers some 2210 km^2 and ranges in elevation from 4500 to 11500 ft asl. Fig. 1.1 shows the layout of the basin with eight smaller sub-basins which have either active or inactive stream gauging stations. The drainage network with gauge sites is shown in fig. 1.2. The relief within the basin is shown in fig 1.3 and the distributions of forest and glacier cover are shown in figs. 1.4 and 1.5. A schematic of the relative positions of the stream gauge network within the major drainage network is shown in fig. 1.6.

A GIS (Terrasoft) has been employed in order to describe basin characteristics and as a basis for further analysis of the data. The superimposition of ***basin and sub-basin boundaries***, the ***relief information*** (derived from digitizing contour lines at 500ft intervals), and the ***land surface types*** (forest, glaciers, lakes and bare rock) allow hypsographic curves for the basins and sub-basins to be derived, illustrated in fig. 1.7. Table 1.1 provides a summary of the areal breakdown of land surface types within sub-basins. It is worth noting that within the basin as a whole there is only some 73 km^2 glacier cover (as calculated from 1950 maps) or 3.3% of the land area. More than half of this glacier area lies within the Lake Louise sub-basin. Table 1.1 also includes data from the adjacent Mistaya basin as the Mistaya has excellent stream gauging records and, because it is very similar in characteristics to the Hector Lake sub-basin, it will be used in later analyses for comparative study.



Fig. 1.1 The Bow River Basin above the Banff stream gauge (BB001), covering 2210 km^2, with sub-basins delineated.



Fig. 1.2 The Bow River Basin above the Banff showing major drainage network and gauge sites for sub-basins.



Fig. 1.3 Relief within the Bow River Basin above the Banff.



Fig. 1.4 Forest distribution within the Bow River Basin above the Banff.



Fig. 1.5 Glacier distribution within the Bow River Basin above the Banff.

BASIN		Total Area	Forest	Glacier	Lake	Rock	%	%	%	%
		km. sq.					Forest	Glacier	Lake	Rock
Bow River at Banff	BB001	2226.66	1160.06	72.86	15.06	978.68	52.10	3.27	0.68	43.95
Lake Louise	BA001	425.55	174.78	45.13	9.04	196.60	41.07	10.61	2.12	46.20
Pipestone River	BA002	307.79	119.71	9.79	0.35	177.94	38.89	3.18	0.11	57.81
Johnson Creek	BA006	124.17	50.94	0.00	0.31	72.92	41.02	0.00	0.25	58.73
Baker Creek	BA007	125.49	40.34	0.09	0.59	84.47	32.15	0.07	0.47	67.31
Hector Lake	BA008	281.41	83.81	33.20	9.04	155.36	29.78	11.80	3.21	55.21
Forty Mile Creek	BB003	133.41	64.95	0.00	0.13	68.33	48.68	0.00	0.10	51.22
Brewster Creek	BB004	107.15	48.73	0.49	0.25	57.68	45.48	0.46	0.23	53.83
Redearth Creek	BB005	154.16	94.12	1.71	1.23	57.10	61.05	1.11	0.80	37.04
Rest of basin		848.97	566.48	15.65	3.18	263.66	66.73	1.84	0.37	31.06
COMPARATIVE BASINS										
Mistaya	DA007	247.21	83.81	26.63	5.84	130.94	33.90	10.80	2.40	52.90
Silverhorn Creek	DA010	19.99	1.91	0.58	0.00	17.50	9.60	2.90	0.00	87.50
Peyto Creek	DA008	22.25	0.00	12.12	0.00	10.13	0.00	54.50	0.00	45.50

Table 1.1 Characteristics of Bow River Basin above Banff and adjacent basins for comparative studies



Fig 1.6 Schematic of relative positions of stream gauges within the Bow River Basin above Banff



Fig. 1.7 Hypsographic curves for the basin of the Bow River above Banff and for the Bow River above Lake Louise

CHAPTER 2 Database II: Streamflow records

A summary of the streamflow records for the Bow River at Banff (BB001) and for 8 sub-basins (BA001,2,6,7,8, BB003,4,5) are presented in this chapter. Fig. 2.1 provides a Time line for the streamflow records for the Bow River above Banff and for gauges within the Mistaya basin contiguous to and immediately north of the Bow basin.

The overall flow record is summarised in fig. 2.2. It is apparent from this figure that there are no very obvious major trends in the 82 period of record. It is clear, however, that there have been extended periods of generally lower than average flows (the 1940s and the last two decades) and periods of generally higher than average flows (the 1960s). Fig. 2.3 further illustrates the discharge records by showing mean monthly flows. Differences in the hydrographs from year to year are apparent and the considerable contribution of the Lake Louise sub-basin is seen in fig 2.4. A further illustration of the dominance of the snowmelt phase in early summer is shown in fig. 2.5.

It will be noted from fig. 2.1 that, fortunately, for the period 1973-76 stream gauging was undertaken at all gauges. This will allow an intercomparison of the records for that time period. Fig. 2.6 shows mean monthly flows at Banff for the 1973-76 period compared to the mean monthly flows for the 82-year period of record. It shows that (again fortunately) within the 4-year 1973-76 record there were higher than average and lower than average years. The hydrographs in this 4-year period showed considerable diversity, as show by the daily records of the Bow River at Banff in fig. 2.7.

To illustrate the differences between sub-basins with different land cover characteristics fig. 2.8 presents mean monthly discharges 1973-76 for the Bow River at Banff, the Bow River below Hector Lake (almost 12% glacier covered) and the Forty Mile Creek (no glacier cover). The data are standardized to eliminate the effect of different areas within the basins. It is apparent that the Hector Lake gauge records high specific yields and that the maximum discharges are delayed until later in the summer.

A data base of discharges within each sub-basin has been linked to the GIS. This allows a mapped description of the relative contributions of sub-basins to overall basin yields. As illustration of this powerful technique, maps of sub-basin yield by month for June to September 1973-76 are presented in the Appendix. The yields have been standardised, ie yields have been expressed as $m^3 s^{-1} km^{-2}$, allowing ready intercomparison of sub-basin yields. This technique could also be applied on a daily or annual basis and will form the basis of future modelling programs.



Fig 2.1 Time line of availability of major series of discharge within the Bow River basin above Banff and in the adjacent Mistaya Basin



Fig. 2.2 Mean annual flows of the Bow river at Banff 1910-1992



Fig. 2.3 Mean monthly discharge of the Bow River at Banff 1909-1992



Fig. 2.4 Mean monthly discharge of the Bow River at Banff and for the Bow River at Lake Louise 1965-1992



Fig. 2.5 Discharge of the Bow River at Banff by month 1910-1992



Fig. 2.6 Mean monthly flows of the Bow River at Banff 1973-76 as a percentage of the mean monthly flows 1911-1992



Fig. 2.7 Bow River at Banff daily discharge 1973-1976



Fig. 2.8 Standardized mean monthly discharges 1973-1976 for the Bow River at Banff, the Bow below Hector Lake and Forty Mile Creek

Chapter 3 Database III: Meteorological records, snowcourse records and records of glacier mass balance

This chapter introduces the meteorological data available within the basin. Essentially only the high quality and consistent records of temperature and precipitation gathered over a long period of time are those from Banff. Great care must be taken when analysing climatic data records. It is known that the Lake Louise records are broken and must therefore be used with caution.

The Banff records are illustrated in fig. 3.1 showing annual temperature and precipitation records from 1895-1992. The temperature record (daily mean values are almost complete during the whole of this period, while the precipitation records are close to complete from 1910 onwards). It is, perhaps possible to detect a slow rise in temperature over this almost 100 year record. There are some periods of higher than average and lower than average temperatures. Of particular note was a marked cool period in the late 1940s. Precipitation shows an annual mean of about 450mm at Banff. The higher mountain areas show much higher mean annual precipitation often in the range 1200 - 1800 mm.

Records of temperature and other elements of the energy balance have been recorded at the Peyto Glacier research facility at 2200 masl, particularly for the period 1966-74. Records of glacier mass balance on Peyto Glacier cover an elevation range of 1000m and are illustrated in summary form in fig. 3.4.These data are not illustrated in detail within this report, but will be used in future analyses, especially for the determination of temperature and precipitation lapse rates, so important for many hydrological models.

Of particular importance for the understanding the hydrology of high mountain areas is the state of the snowpacks immediately prior to spring melt. Heavy winter snowpacks will produce increased runoff especially during the early summer period and may well decrease the contribution of glacier melt waters during the entire summer. Conversely, low winter snowpacks can lead to reduced overall summer flows, but can increase the yield and the relative importance of glacier ice melt.

Thus emphasis has been placed on the compilation of the snow course and the snow pillow data within the basin. A Time line for availability of snowcourse data is found in fig. 3.2. Table 3.1 and fig. 3.3 illustrate the April 1st records at some of the courses with long records. It will be noted from these summary presentations that there is a very strong covariance between the courses indicating a uniformity of snow inputs over the whole basin. This is an important finding as it will influence the types of basin modelling to be employed in future analyses.

Cautionary note: it was found that in several instances the computer files containing summary data were in error; sometimes errors were very great which could easily lead to erroneous analyses; these errors have been corrected as far as possible by reference to primary data files, but all data files will have to be rechecked for accuracy and consistency.



Fig. 3.1 Annual temperature and precipitation records at Banff, 1895-1992



(a)



(b)

Fig. 3.2(a) Timeline of data availability and **(b)** location of selected snow courses

Snow course data 1937 - 1989 for April 1 surveys

	Bow River (1580m)	Chateau Lawn (1740m)	Mirror Lake (2030m)	Pipestone Upper (1615m)		Bow River (1580m)	Chateau Lawn (1740m)	Mirror Lake (2030m)	Pipestone Upper (1615m)	Bow Summit (old) (2080m)	Bow Summit (new) (2080m)	Bow Summit (composite) (2080m)
1937	97			117	1964	137	178	193	152			
1938	208			282	1965	284	323	378	259			
1939	180			183	1966	213	226	274	193			
1940	132	142	168	142	1967	437	478	538	434			
1941	135	135	203	150	1968	170	234	254	163	328		458
1942	114	173	180	117	1969	196	310	320	208			
1943	168	178	224	160	1970	119	152	163	109	127		177
1944	137	157	178	137	1971	259	356	368	251	323		451
1945	104	173	160	127	1972	386	480	561	406	546		763
1946	216	277	353	234	1973	175	325	279	150	300		419
1947	241	282	358	246	1974	277	381	419	193	533		745
1948	229	292	310	234	1975	203	257	236	190	264		369
1949	160	239	211	147	1976	269		361	251	460		643
1950	274	384	427	292	1977	147		175	114	170		237
1951	274	389	460	338	1978	198	216	244	175	310		433
1952	229	262	315	226	1979	191	259	234	198	251	361	361
1953	213	300	274	218	1980	232	333	320	234	320	434	434
1954	330	391	452	218	1981	244	328	318	246		381	381
1955	163	229	239	160	1982	234	300	295	284		353	353
1956	323	373	427	310	1983	206	277	284	229		340	340
1957	269	307	378	264	1984	142	196	239	152		330	330
1958	218	267	297	216	1985	175	211	211	196		315	315
1959	259	305	310	264	1986	236	290	345	287		439	439
1960	150	160	262	145	1987	218	279	302	221		302	302
1961	224	310	333	236	1988	160	213	251	191		320	320
1962	206	264	279	198	1989	196	292	284	234		384	384
1963	147	188	218	152								

Table 3.1 April 1st water equivalents (mm) for selected snowcourses 1937-1989



Fig. 3.3 April 1st water equivalents (mm) for the four snowcourses with the longest records



Fig. 3.4 Peyto Glacier mass balance 1966-84, with extrapolations back to 1950

Chapter 4 Database IV: Glacier inventory and selected coverage by aerial photography

It had been hoped that considerably more progress in compiling data from the Canadian Glacier Inventory would have been achieved. Unfortunately the data archives within the National Hydrology Research Institute could not readily be accessed. Illustration of the glacier coverage within the Bow basin as defined and referenced within the Glacier Inventory is given in fig. 4.1.

Data for the glacier inventory for the Waputik Mountain area (within the western section of the Lake Louise sub-basin) are available from Müller et al 1967. The map from this publication is given in fig. 4.2 and summary statistics from the publication are given in table 4.1. It will be noted that the glacier reference numbers within the maps in figs. 4.1 and 4.2 are not consistent. This aspect of the glacier inventory work needs to be rectified before future analyses may be undertaken.

Over the last few decades many surveys using air photography have been conducted. Some of these surveys are illustrated in figs 4.2(a,b and c).



Fig. 4.1 Glacier inventory base map of the Upper Bow River and adjacent basins from the Glacier Atlas of Canada



Fig. 4.2 Map of the Glacier Inventory of the Waputik Mountains from Müller et al 1967

Bow River Headwaters

Ident	No.	Longitude	Latitude	U.T.M	Orientation AC	Orientation AB	Elevations High	Elevations L.Ex	Elevations L.Tl	Snow	Date	Mean Elevation Accumulation	Mean Elevation Ablation
5BAC	2	w 116 20.0	n 51 28.0	11UNH546257021	N	N	2347	2134	2134		22 08 66	2225	
5BAC	3	w 116 23.0	n 51 30.5	11UNH543157063	S	SE	2774	2347	2347	2499	22 08 66	2652	2499
5BAC	4	w 116 22.5	n 51 33.0	11UNH543557126	SE	SE	2713	2225	2225	2469	22 08 66	2530	2347
5BAC	5	w 116 18.5	n 51 29.5	11UNH547857047	N	N	2605	2499	2499		22 08 66	2591	
5BAC	6	w 116 18.5	n 51 30.0	11UNH547857053	E	E	2377	2195	2195		22 08 66	2347	
5BAC	7	w 116 19.0	n 51 30.5	11UNH547757064	NE	E	2225	2103	2103		22 08 66	2195	
5BAD	1	w 116 22.5	n 51 34.0	11UNH543357126	N	NW	2804	2225	2225		22 08 66	2682	
5BAD	2	w 116 23.5	n 51 34.0	11UNH542357123	N	NW	2774	2347	2347		22 08 66	2652	
5BAD	3	w 116 25.0	n 51 34.0	11UNH540257126	N	NW	3048	2134	2134	2438	22 08 66	2652	2286
5BAD	4	w 116 27.5	n 51 34.5	11UNH537857141	N	NE	3048	2286	2286	2530	22 08 66	2896	2530
5BAD	5	w 116 27.5	n 51 35.5	11UNH537457160	SE	SE	2830	2330	2330	2530	22 08 66	2713	2499
5BAD	6	w 116 27.5	n 51 37.0	11UNH537957187	SW	SE	2865	2445	2445	2591	22 08 66	2743	2530
5BAD	7	w 116 25.5	n 51 37.5	11UNH539757193	S	S	2743	2575	2575		22 08 66	2698	
5BAD	8	w 116 26.0	n 51 38.0	11UNH540857200	NE	NE	2377	2316	2316		22 08 66	2347	
5BAD	9	w 116 26.0	n 51 38.0	11UNH539457202	NW	N	2743	2408	2408	2438	22 08 66	2682	2499
5BAD	10	w 116 27.0	n 51 38.5	11UNH538057217	SE	E	2438	2286	2286	2347	22 08 66	2408	2316
5BAD	11	w 116 27.5	n 51 38.0	11UNH537757202	NW	NW	2896	2575	2575		22 08 66	2790	
5BAD	12	w 116 27.5	n 51 37.5	11UNH537257194	NW	NW	2743	2682	2682		22 08 66	2713	
5BAD	13	w 116 30.0	n 51 38.0	11UNH534657203	N	N	2926	2469	2469	2560	22 08 66	2665	2530
5BAD	14	w 116 28.0	n 51 37.5	11UNH535757198	NW	NW	2591	2452	2452	2499	22 08 66	2499	2499
5BAD	15	w 116 29.0	n 51 38.0	11UNH535857198	N	N	2438	2256	2256	2316	22 08 66	2377	2286
5BAD	16	w 116 30.5	n 51 39.0	11UNH533957216	NE	NE	2774	2317	2317	2560	22 08 66	2652	2484
5BAD	17	w 116 31.0	n 51 39.5	11UNH533457233	E	E	2804	2652	2652		22 08 66	2743	
5BAD	18	w 116 31.5	n 51 40.0	11UNH533257240	E	E	2896	2713	2713		22 08 66	2758	
5BAD	19	w 116 31.0	n 51 40.5	11UNH533657246	SE	SE	2743	2576	2576	2606	22 08 66	2652	2591
5BAD	20	w 116 30.0	n 51 41.5	11UNH535657264	NE	NE	2591	2469	2469		22 08 66	2560	
5BAD	21	w 116 30.0	n 51 42.0	11UNH534557276	N	N	2499	2377	2377		22 08 66	2438	

Bow River Headwaters

Ident	No.	Total Length			Width	Surface Area					AAR	Depth	Volume **		Type
		Ablation	Exp	Total		Exp	Total		Ablation						
5BAC	2		0.4	0.4		0.34	0.34	2			100	10	0.003	3	600010
5BAC	3	0.2	3.1	3.1	1.5	4.3	4.3	2	0.34	2	92	50	0.215	3	600010
5BAC	4	1.3	3.2	3.2	0.8	2.9	2.9	2	0.59	2	79	50	0.145	4	520012
5BAC	5		0.3	0.3		0.06	0.06	2			100	10	0.001	4	600010
5BAC	6		0.4	0.4	0.2	0.08	0.08	2			100	10	0.001	4	600000
5BAC	7		0.4	0.4	0.2	0.05	0.05	2			100	10	0.001	4	640010
5BAD	1		0.6	0.6	0.4	0.27	0.27	2			100	10	0.003	4	640402
5BAD	2		0.8	0.8	0.4	0.31	0.31	2			100	10	0.003	4	603312
5BAD	3	1.5	3.9	3.9	1.2	5.9	5.9	2	1.7	2	71	70	0.413	3	520412
5BAD	4	0.5	2.7	2.7		2.9	2.9	2	0.8	2	74	70	0.203	3	503311
5BAD	5	1.3	3.2	3.2	1	4.9	4.9	2	1	2	79	70	0.343	4	503012
5BAD	6	1.1	1.9	1.9	0.8	1.5	1.5	2	0.44	2	71	50	0.008	4	530010
5BAD	7		0.5	0.5	0.2	0.25	0.25	2			100	10	0.002	4	600012
5BAD	8		0.4	0.4		0.12	0.12	2			100	10	0.001	4	600010
5BAD	9	0.1	1.6	1.6	0.7	1.5	1.5	2	0.06	2	66	25	0.004	4	600212
5BAD	10	0.1	0.6	0.6	0.2	0.14	0.14	2			100	10	0.001	4	630000
5BAD	11		1	1	0.7	0.45	0.45	2			100	10	0.004	4	600010
5BAD	12		0.3	0.3	0.2	0.06	0.06	2			100	10	0.001	4	600310
5BAD	13	0.3	2.6	2.6	0.7	2	2	2	0.09	2	95	25	0.05	4	600210
5BAD	14	0.1	0.4	0.4		0.11	0.11	2	0.02	2	80	10	0.001	4	640002
5BAD	15	0.2	0.5	0.5		0.27	0.27	2	0.06	2	78	10	0.003	4	640002
5BAD	16	1.6	3.1	3.1	0.5	5.1	5.1	2	0.52	2	90	100	0.51	3	523411
5BAD	17		0.2	0.2		0.06	0.06	2			100	10	0.001	4	640000
5BAD	18		0.2	0.2		0.2	0.2	2			100	10	0.002	4	630010
5BAD	19	0.1	0.7	0.7		0.38	0.38	2	0.08	2	79	10	0.004	4	630002
5BAD	20		0.5	0.5		0.19	0.19	2			100	10	0.002	4	640000
5BAD	21		0.3	0.3		0.6	0.6	2			100	10	0.001	4	600012
		6.9	26		7	27.21			4.77				1.56		

** Note - For volume accuracy rating "4" read "3" and for "3" read "2"

Summary of the glacier inventory of the Bow River headwaters of the Waputik Mountains

Location	No.	Volume	Area	Max. size	Elevation		Transient snow line	
		cubic k	square km	square km	Highest	Terminus	From	To
5Bac	6	0.366	7.73	4.3	2800	2100	2450	
5Bad	21	1.56	27.21	5.9	2900	2100	2350	2600

Table 4.1 Summary data from glacier inventory of the Waputik Mountains from Müller et al 1967

Air Photographic Coverage of the Bow River Basin

Table 4.2(a) Air Photographs from 1951 - 1952 Scale 1:40 000

Flight Line	Year flown	Photograph Numbers
AS162-5110	1951	124-136
AS162-5110	1951	195-208
AS163-5109	1951	54-66
AS163-5108/09	1951	20-33
AS164-5108	1951	140-155
AS164-5107	1951	5-11
AS165-5106/07	1951	224-235
AS165-5106	1951	78-90
AS166-5105	1951	174-185
AS166-5104	1951	38-50
AS166-5103/04	1951	14-25
AS167-5103	1951	152-158
AS167-5102	1951	26-30
AS168-5101	1951	128-135
AS168-5100	1951	1-6
AS169-5200	1952	104-110

Table 4.2(b) Air Photographs from the year 1977 July / August, Scale 1:50 000

Flight Number	Photograph Numbers
A24783	55-79
A24783	71-85
A24783	95-110
A24776	134-140
A24776	65-70
A24805	105-115
A24805	55-61
A24739	100-105
A24783	125-136

Table 4.2(c) Air Photographs from the year 1993,September, Scale 1:50 000

Flight Number	Photograph Numbers	
A27991	94-104	
A27991	10-19	
A27988	284-293	Partial Cloud
A27988	262-255	259-260 Cloud
A27988	205-216	209-210 Cloud
A27988	118-129	

Bibliography

Glacier Studies on the Eastern Slopes of the Rocky Mountains

BINDA, G.G., 1984. Fluvioglacial sediment and hydro chemical dynamics, Peyto Glacier, Alberta. M.A. thesis, Department of Geography, University of Ottawa, Ottawa, Ontario, 139 pp.

BINDA, G.G., JOHNSON, P.G. and POWER, J.M., 1985. Glacier control of suspended-sediment and solute loads in a Rocky Mountain basin. In - Water Quality Evolution within the Hydrological Cycle of Watersheds, Proceedings of the Canadian Hydrology Symposium No.15 - 1984, 10-12 June 1984, Quebec, NRCC No.24633, Associate Committee on Hydrology, National Research Council of Canada, Volume I, p 309-327.

BRADLEY, C., 1990. Water quality dynamics in the meltwaters draining Peyto Glacier, Alberta. M.A. thesis, Department of Geography, Wilfrid Laurier University, Waterloo, Ontario, 133 pp.

BRUGMAN, M.M., 1992. Search for trends in glacier mass balance from western Canada. In - Using Hydrometric Data to Detect & Monitor Climatic Change, G.W. Kite and K.D. Harvey (Editors), NHRI Symposium No.8, Proceedings of NHRI Workshop No. 8, 8-9 April 1992, Saskatoon, Saskatchewan, National Hydrology Research Institute, Saskatoon, Saskatchewan and Surveys and Information Systems Branch, Ottawa, Ontario, p 233-244.

BRUGMAN, M.M. and LI, L., 1992. Peyto Glacier DEM and SAR image simulation project: Generation of slope and aspect grids from digital elevation model and a first simulation of the NASA/JPL SAR image taken on May 3, 1991. NHRI Internal Planning Report No. 92002, Hydrological Sciences Division, National Hydrology Research Institute, Environment Canada, Saskatoon, Saskatchewan, 25 March, NHRI Internal Planning Report No. 92002.18 pp.

BRUNGER, A.G., 1966. The Drummond, Hector and Peyto glaciers - their wastage and deposits. M.Sc. thesis, Department of Geography, University of Alberta, Edmonton, Alberta, 100 pp.

BRUNGER, A.G., NELSON, J.G. and ASHWELL, I.Y., 1967. Recession of the Hector and Peyto glaciers: further studies on the Drummond Glacier, Red Deer Valley area, Alberta. Canadian Geographer, 11(1), p 35-48

CAMPBELL, P.I., REID, I.A., and SHASTAL, J., 1969. Glacier Survey in Alberta. Water Survey of Canada, Inland Waters Branch, Department of Energy, Mines and Resources. Report No.4. Ottawa. 15 pp.

CANADA, ENVIRONMENT CANADA, 1976. Canadian participation in the International Hydrological Decade. Final Report. Volume 3 - Research basins. Canadian National Committee for the International Hydrological Decade, Inland Waters Directorate, Environment Canada, Ottawa, Ontario, 184 pp.

CANADA, FISHERIES AND ENVIRONMENT CANADA, 1977. Canadian participation in the International Hydrological Decade. Final report: Volume 1 - summary of activities. Canadian National Committee for the International Hydrological Decade, Inland Waters Directorate, Fisheries and Environment Canada, Ottawa, Ontario, Volume 1, 79 pp.

CARTER, R. L., 1954. Study of Glaciers on Eastern Slope of the Rocky Mountains in Banff and Jasper National Parks, 1952 and 1954. Report No.2160-(52 & 54),surveys under the supervision of R.D. May, Dominion Water and Power Bureau, Forests and Scientific Services Branch, Department of Mines and Resources, Calgary, Alberta. 32 pp.

CARTER, R.L., BLAIR, R.V., FOWLER, E.D., and THORSON, K.M., 1956. Study of Glaciers in Banff and Jasper National Parks, 1956. Report No.2160-(56), Water Resources Branch, Department of Northern Affairs and National Resources, Calgary, Alberta. 27 pp.

CAUTLEY, R.W., and WHEELER, A.O., 1924. Interprovincial Boundary Commission, Boundary between Alberta and British Columbia. Scale 1:62500, maps 13-21, based on surveys from 1917. Office of the Surveyor General, Ottawa, Ontario.

CAUTLEY, R.W., and WHEELER, A.O., 1924. Report of the Commission Appointed to Delimit the boundary between the Provinces of Alberta and British Columbia, Part II 1917-1921, from Kicking Horse Pass to Yellowhead Pass. F.A.Aclund Printing. Ottawa. 157 pp.

CHAPMAN, E.F., MORTON, G.H., THORSON, K.M., McGEACHY, D.A., ELDER, V.S. and LOEPPKY, K.D., 1960. Survey of Glaciers on the Eastern Slopes of the Rocky Mountains in Banff and Jasper National Parks, 1960. Report No.2160-(60),Water Resources Branch, Department of Northern Affairs and National Resources, Calgary, Alberta. 31 pp.

COLLIER, E.P., 1957. Glacier Variation and Trends in Run-off in the Canadian Cordillera. Comptes Rendus et Rapports: Assemblee Generale de Toronto, 1957,Tome iv,(1957). p 344-357.

COLLINS, D.N., 1984. Water and mass balance measurements in glacierised drainage basins. Geografiska Annaler, 66A(3), p 197-214.

CHURCH, M. and GILBERT, R., 1975. Proglacial fluvial and lacustrine environments. In - Glaciofluvial and Glaciolacustrine Sedimentation, A.V. Jopling and B.C. MacDonald (Editors), Society of Economic Paleontologists and Mineralogists Special Publication No.23, Tulsa, Oklahoma, October, p 22-100.

DALTON, A., 1991. Peyto Glacier mass balance 1985-1990. NHRI Internal Planning Report No.91001 [limited distribution], Hydrological Sciences Division, National Hydrology Research Institute, Environment Canada, Saskatoon, Saskatchewan, NHRI Internal Planning Report No.91001.June, 38 pp.

DAVIES, K.F., 1964. Survey of glaciers on eastern slope of Rocky Mountains in Banff and Jasper National Parks, 1964. Report No.2160-(64), Water Resources Branch, Department of Northern Affairs and National Resources, Calgary, Alberta, mimeo, 21 pp.

DAVIES, K.F., WARNER, L.A., ANDERSON,, J.E. and DAHL, B.M., 1970. Survey of glaciers on the eastern slopes of the Rocky Mountains in Banff and Jasper National Parks - 1970 report. Internal Report, Water Survey of Canada Division, Inland Waters Branch, Department of Energy, Mines and Resources, Calgary, Alberta, 29 pp.

DAVIS, D.A., LOEPPKY, K.D., SAPP, E. and ADAMS, R.E., 1962. Study of glaciers on eastern slope of Rocky Mountains in Banff and Jasper National Parks, 1962. Report No.2160-(62), Water Resources Branch, Department of Northern Affairs and National Resources, Calgary, Alberta, October, mimeo, 32 pp.

DAVIES, D.A., LOEPPKY, K.D., SAPP, E., and ADAMS, R.E., 1962. Survey of Glaciers on the Eastern Slopes of the Rocky Mountains In Banff and Jasper National Parks, 1962. Report No.2160-(62).Water Resources Branch, Department of Northern Affairs and National Resources, Calgary, Alberta. pp 32.

DAVIES, K.F., FROELICH, C.R., HEINZE, L., and KERBER, R., 1966. Survey of Glaciers on the Eastern Slopes of the Rocky Mountains in Banff and Jasper National Parks, 1966. Report No.2160-(66),Water Resources Branch, Department of Northern Affairs and National Resources, Calgary, Alberta. 30 pp.

DERIKX, L., 1972. Hydrological characteristics of Peyto Glacier. In - Guidebook to the International Symposia on the Role of Snow and Ice in Hydrology, Banff, Alberta, 1972, Canadian National Committee for the I.H.D., p 79-84.

DERIKX, L. and LOIJENS, H., 1970. Model of runoff from glaciers. Hydrology Symposium No.8, Subcommittee on Hydrology, Associate Committee on Geodesy and Geophysics, National Research Council, Vol.1, p 153-199.

FIELD, W.O., Jr and HEUSSER, C.J., 1954. Glacier and botanical studies in the Canadian Rockies, 1953. Canadian Alpine Journal, Vol.37, p 127-140.

FOESSEL,D.G., 1974. An analysis of the temperature distribution over the Peyto Glacier, Alberta. M.Sc. thesis, Department of Geography, Guelph University, Guelph, Ontario, 137 pp.

FOUNTAIN, A.G. and TANGBORN, W., 1985. Overview of contemporary techniques. Techniques for prediction of runoff from glacierized areas, G.J. Young (Editor), IAHS Publication No.149, International Association of Hydrological Sciences, p 27-41.

FOWLER, E.D., CHAPMAN, E.F., THORSON, K.M., and RAINSBERRY, K.C., 1958. Study of Glaciers in Banff and Jasper National Parks,1958. Report No.2160-(58). Water Resources Branch, Department of Northern Affairs and National Resources, Calgary, Alberta. 29 pp.

GADD, B., 1988. Handbook of the Canadian Rockies. Corax Press. Alberta. 878 pp.

GLENDAY, P.J., 1991. Determination of morphologic and volumetric change, Peyto Glacier, Alberta, 1966-1989. M.A. thesis, Department of Geography, Wilfrid Laurier University, Waterloo, Ontario, 130 pp.

GLOSSOP, J., MORTON, G.H., ANDERSON, J.E., SLOBOSZ, F. and CLAYTON, V., 1968. Survey of glaciers on the eastern slope of the Rocky Mountains in Banff and Jasper National Parks, 1968. Report No.2160-(68), Water Survey of Canada, Inland Waters Branch, Department of Energy, Mines and Resources, Calgary, Alberta, mimeo, 29 pp.

GOODISON, B., 1969. The distribution of global radiation over Peyto Glacier and a statistical runoff model for Peyto Glacier. In - Proceedings of the North Saskatchewan Headwaters meeting, 5 December 1969, Ottawa, Inland Waters Branch, Department of Energy, Mines and Resources, Ottawa, Ontario, mimeo, unpaginated, 2 pp.

GOODISON, B., 1969. A preliminary investigation of the distribution of global radiation over Peyto Glacier. M.A. thesis, Department of Geography, University of Toronto.

GOODISON, B., 1970. The relation between ablation and global radiation over Peyto Glacier, Alberta. In - Glaciers, Proceedings of IHD Workshop Seminar 1970, Vancouver, B.C., Canadian National Committee for the International Hydrological Decade, p 39-42.

GOODISON, B., 1972a. An analysis of climate and runoff events for Peyto Glacier, Alberta. Inland Waters Directorate Scientific Series No.21, Water Resources Branch, Environment Canada, Ottawa, Ontario, 29 pp.

GOODISON, B., 1972b. The distribution of global radiation over Peyto Glacier, Alberta. Inland Waters Directorate Scientific Series No.22, Water Resources Branch, Environment Canada, Ottawa, Ontario, 22 pp.

HENOCH, W.E.S., 1969. Estimate of glaciers' volumetric change and its contribution to the discharge of the North Saskatchewan River (Banff National Park, Alberta) 1948- 1966. In - Proceedings of the North Saskatchewan Headwaters meeting, 5 December 1969, Ottawa, Inland Waters Branch, Department of Energy, Mines and Resources, Ottawa, Ontario, mimeo, unpaginated, 3 pp.

HENOCH, W.E.S., 1971. Estimate of glaciers' secular (1948-1966) volumetric change and its contribution to the discharge in the upper North Saskatchewan River basin. Journal of Hydrology, Vol.12, p 145-160.

HENOCH, W.E.S. and CROIZET, J., 1976. The Peyto Glacier map/a three-dimensional depiction of mountain relief. The Canadian Cartographer, 13(1), June, p 69-86.

HEUSSER, C.J., 1954. Glacier fluctuations in the Canadian Rockies. IASH Publication No.39, p 493-497.

HEUSSER, C.J., 1956. Postglacial Environments in the Canadian Rocky Mountains. Ecological Monographs, Vol.26, No.4. p 263-302.

HISOP, R.W.P., 1974. Stable Isotopes for identifying accumulation and ablation processes on the Wapta Icefield. M.Sc. Thesis, University of Calgary. 87 pp.

HOBSON, G.D. and JOBIN, C., 1975. A seismic investigation - Peyto Glacier, Banff National Park and Woolsey Glacier, Mount Revelstoke National Park. Geoexploration, 13(2), April, p 117-127.

HOLDSWORTH, G., and DEMUTH, M.N., 1987. Annex 3: Mass Balance - Peyto Glacier. In - National Hydrology Research Institute internal review - glaciology programme: (includes snow but excludes ice in rivers), T.M. Dick (editor), NHRI Internal Planning Report No.87003e, Surface Water Division, National Hydrology Research Institute, Environment Canada, Saskatoon, Saskatchewan, 18 December, NHRI Internal Planning Report No.87003e. 2 pp.

HOLDSWORTH, G., POWER, J. and CHRISTIE, R., 1983. Radar ice thickness measurements on Peyto Glacier. Internal report, Snow and Ice Division, National Hydrology Research Institute, Environment Canada, Ottawa, Ontario, December, 18 pp.

JOHNSON, P.G., 1982. Prediction of runoff from glacierized areas: Canada. Report to Working Group on the Prediction of Glacier Runoff, International Commission on Snow and Ice, from Department of Geography, University of Ottawa, Ottawa, Ontario, 16 pp.

JOHNSON, P.G., 1985. The importance of glacier mass balance programmes in Canada. Unpublished review paper for Subcommittee on Glaciers, Department of Geography, University of Ottawa, Ottawa, Ontario, 10 pp.

JOHNSON, P.G. and POWER, J.M., 1985. Flood and landslide events, Peyto Glacier terminus, Alberta, Canada, 11-14 July 1983. Journal of Glaciology, 31(108), p 86-91.

KENNEDY, S.K., 1975. Sedimentation in a glacier-fed lake. M.Sc. thesis, Department of Geological Sciences, University of Illinois at Chicago Circle, Chicago, Illinois, 55 pp.

LEONARD, E.M., 1981. Glaciolacustrine sedimentation and Holocene glacial history, northern Banff National Park, Alberta. Ph.D. thesis, University of Colorado, Boulder, Colorado, 271 pp.

LEONARD, E.M., 1985. Glaciological and climatic controls on lake sedimentation, Canadian Rocky Mountains. In - Climate and Palaeoclimate of Lakes, Rivers and Glaciers, M. Kuhn (Editor), Proceedings of Symposium, 4-7 June 1984, Igls, Austria, Zeitschrift fur Gletscherkunde und Glazialgeologie, Heft 21, p 35-42.

LEONARD, E.M., 1986. Use of lacustrine sedimentary sequences as indicators of Holocene glacial history, Banff National Park, Alberta, Canada. Quaternary Research, 26(2), p 218-231.

LEONARD, E.M., 1986. Varve studies at Hector Lake, Alberta, Canada, and the relationship between glacial activity and sedimentation. Quaternary Research, 25(2), p 199-214.

LOUENS, H.S., 1969. Runoff from the headwaters area of the North Saskatchewan River. In - Proceedings of the North Saskatchewan Headwaters meeting, 5 December 1969, Ottawa, Inland Waters Branch, Department of Energy, Mines and Resources, Ottawa, Ontario, mimeo, unpaginated, 2 pp.

LOUENS, H.S., 1970a. Assessment of glacier melt contribution and stream flow hydrograph synthesis in the North Saskatchewan headwaters, Alberta. Internal report, Inland Waters Directorate, Department of the Environment, Ottawa, Ontario, 37 pp.

LOUENS, H.S., 1970b. Glacier contribution to the flow of the North Saskatchewan River at Saskatchewan Crossing, Rocky Mountains, Alberta. Paper presented at Workshop Seminar on "Evaluation of Basin Models for Alpine Areas", 27-28 July 1970, Wapta Lodge, Field, B.C., Abstract, 6 pp.

LOUENS, H.S., 1974. Streamflow formation in the Mistaya river basin, Rocky Mountains, Canada. Western Snow Conference, Proceedings, 42nd Annual Meeting, 16-20 April 1974, Anchorage, Alaska, p 86-95.

LØKEN, O.H., 1970. Glacier studies in the Canadian IHD program. In - Glaciers, Proceedings of IHD Workshop Seminar 1970, Vancouver, B.C., Canadian National Committee for the International Hydrological Decade, p 1-4.

MARTZ, L., and LI, L., 1992. Digital elevation model generation, Peyto Glacier area: project summary. NHRI Contract report No. 92001, Contract No.KW504-1-1419, Department of Geography, University of Saskatchewan, Saskatoon, Saskatchewan, Hydrological Sciences Division, National Hydrology Research Institute, Environment Canada, Saskatoon, Saskatchewan, CR-92001. 10 pp.

MAY, R.D., BLAIR, R.V., and HARRY, R.M., 1950. Survey of Glaciers on the Eastern Slopes of the Rocky Mountains in Banff and Jasper National Parks,1950. Report No.2160-(50). Dominion Water and Power Bureau, Surveys and Engineering Branch, Department of Mines and resources, Calgary, Alberta. 34 pp.

McCARTHY, D.P., 1985. Dating Holocene geomorphic activity of selected landforms in the Geikie Creek Valley, Mount Robson Provincial Park, British Columbia. M.Sc. thesis, Department of Geography, University of Western Ontario, London, Ontario, September, 303 pp.

McFARLANE, W.T., 1945. Study of glaciers in Banff, Yoho and Jasper National Parks, 1945. Report No.2160-(45), Dominion Water and Power Bureau, Surveys and Engineering Branch, Department of Mines and Resources, Vancouver, B.C., manuscript, 23 pp.

McFARLANE, W.T., 1946a. Glacier Investigations in Banff, Yoho and Jasper National Parks. Canadian Alpine Journal. Vol29, No.2. p 265-273.

McFARLANE, W.T., 1946b. Study of Glaciers in Banff and Jasper National Parks, 1946. Report No.2160-(46). Dominion Water and Power Bureau, Surveys and Engineering Branch, Department of Mines and Resources, Vancouver, British Columbia. 34 pp.

McFARLANE, W.T., 1947. Study of Glaciers in Banff and Jasper National Parks, 1947. Report No.2160-(47). Dominion Water and Power Bureau, Surveys and Engineering Branch, Department of Mines and Resources, Vancouver, British Columbia. 40 pp.

McFARLANE, W.T., and MAY, R.D. 1948. Study of Glaciers in Banff and Jasper National Parks, 1948. Report No.2160-(48). Dominion Water and Power Bureau, Surveys and Engineering Branch, Department of Mines and Resources, Calgary, Alberta. 32 pp.

McFARLANE, W.T., BLAIR, R.V. and OZGA, W.J., 1950. Glacier survey in Banff and Jasper National Parks, 1949. Canadian Alpine Journal, Vol.33, p 120-126.

MEEK, V., 1948. Glacier Observations in the Canadian Cordillera. Canadian Geographical Journal. Vol 37, No.5. p 190-209.

MERCER, J.H., 1958. Chapter 2: Glaciers of the Canadian Rocky Mountains. In - Geographic Study of Mountain Glaciation in the Northern Hemisphere (Contract No. DA19-129-QM-409), Department of Exploration and Field Research, American Geographical Society, New York, June, 23 pp.

MOKIEVSKY-ZUBOK, O., OMMANNEY, C.S.L. and POWER, J., 1985. NHRI glacier mass balance, 1964-1984 (Cordillera and Arctic). NHRI internal report, Glacier Section, Surface Water Division, National Hydrology Research Institute, Environment Canada, Ottawa, Ontario, November, 59 pp.

MÜLLER, F., OMMANNEY, C.S.L. and STANLEY, A.D., 1967. Three pilot studies for the IHD world inventory of glaciers. Abstracts of Papers, Volume VI, General Assembly of Berne, International Association of Scientific Hydrology, International Union of Geodesy and Geophysics, 33 pp.

MUNRO, D.S., 1975. Energy exchange on a melting glacier. Ph.D. thesis, Department of Geography, McMaster University, Hamilton, Ontario, 182 pp.

MUNRO, D.S. and YOUNG, G.J., 1982. An operational net shortwave radiation model for glacier basins. (Paper No.1W1909) Water Resources Research, 18(2), April, p 220-230.

NELSON, J,G., ASHWELL, I.Y., and BRUNGER, A.G. 1966. Recession of the Drummond Glacier, Alberta. Canadian Geographer. Vol10, No2, p71-81.

OMMANNEY, C.S.L., 1969. Glacier inventory of the Rocky Mountains. In - Proceedings of the North Saskatchewan Headwaters meeting, 5 December 1969, Ottawa, Inland Waters Branch, Department of Energy, Mines and Resources, Ottawa, Ontario, mimeo, unpaginated, 4 pp.

OMMANNEY, C.S.L., 1971. Glacier Surveys by the District Personnel of the Water Survey of Canada. Glacier Inventory Note #1. The Victoria Glacier, Canada. Glaciology Division, Inland Waters directorate. Water Resources Branch. Department of the Environment, Ottawa Ontario, 18 pp.

OMMANNEY, C.S.L., 1972a. Glacier inventory. In - Guidebook to the International Symposia on the Role of Snow and Ice in Hydrology, Banff, Alberta, 1972, Canadian National Committee for the I.H.D., p 84-87.

OMMANNEY, C.S.L., 1972b. Glacier surveys by District personnel of the Water Survey of Canada: 2. Peyto Glacier. Glacier Inventory Note No.7, Glaciology Division, Water Resources Branch, Inland Waters Directorate, Department of the Environment, Ottawa, Ontario, June, 20 pp.

OMMANNEY, C.S.L., 1976. Canadian glacier variations and mass balance changes 1970- 1974. Quadrennial report to the Permanent Service on the Fluctuation of Glaciers, Glaciology Division, Environment Canada, Ottawa, Ontario, 18 pp.

OMMANNEY, C.S.L., 1977. Peyto Glacier map: 1:10'000. In - Fluctuations of glaciers 1970-1975 (Vol.III): a contribution to the International Hydrological Programme, Compiled for the Permanent Service on the Fluctuations of Glaciers of the IUGG-FAGS/ICSU by F. Muller, International Commission of Snow and Ice of the International Association of Hydrological Sciences, UNESCO, 259 pp.

OMMANNEY, C.S.L., 1987a. Peyto Glacier: a compendium of information prepared for parks Canada. NHRI Contribution No.87062, Surface Water Division, National Hydrology Research Institute, Environment Canada, Saskatoon, Saskatchewan, NHRI Contribution No. CS-87062. 45 pp.

OMMANNEY, C.S.L., 1990. Peyto Glacier. In - Extract from glaciers of the Canadian Rockies, contribution to the satellite Image Atlas of Glaciers to be published by the U.S. Geological Survey, NHRI Contribution No.90059, National Hydrology Research Institute, Environment Canada, Saskatoon, Saskatchewan, NHRI Contribution No. CS-90059. 13 pp.

OMMANNEY, C.S.L., 1991a. Glacier monitoring: international perspective. NHRI Contribution No.91006, Report presented at the U.S. Parks Service Workshop on Glacier Monitoring, Birchwood Camp, Eagle River, Alaska, 5-6 February 1991, National Hydrology Research Institute, Environment Canada, Saskatoon, Saskatchewan, February,NHRI Contribution No. CS-91006. 71 pp.

OMMANNEY, C.S.L., 1992a. Canadian glacier references (1991-1992) and recent work. NHRI Contribution No.91039, Hydrological Sciences Division, National Hydrology Research Institute, Environment Canada, Saskatoon, Saskatchewan, July, NHRI Contribution No.CS-91039. 35 pp.

ØSTREM, G., 1966. Mass balance studies on glaciers in western Canada, 1965. Geographical Bulletin, 8(1), p 81-107.

ØSTREM, G., 1973. Runoff forecasts for highly glacierized basins. IAHS Publication No.107, p 1111- 1132.

POWER, J.M. and YOUNG, G.J., 1979. Application of the UBC watershed model to Peyto Glacier basin. Canadian Hydrology Symposium: 79 - Cold Climate Hydrology, Proceedings, 10-11 May 1979, Vancouver, B.C.,

Associate Committee on Hydrology, National Research Council of Canada (NRCC 17834), Ottawa, Ontario, p 217-228.

REID, I.A., 1972. Glacier surveys by the Water Survey of Canada. Role of Snow and Ice in Hydrology; Proceedings of the Banff Symposium, Sept.1972, Vol 2, No 107. IAHS, Geneva, Switzerland, WMO-IAHS-UNESCO, p 1133-1143.

REID, I.A. and CHARBONNEAU, J.O.G., 1972. Glacier surveys in Alberta. Inland Waters Directorate Report Series No.22, Department of the Environment, Ottawa, Ontario, 17 pp.

REID, I.A. and CHARBONNEAU, J.O.G., 1978. Glacier surveys in Alberta - 1975. Inland Waters Directorate Report Series No.60, Water Resources Branch, Fisheries and Environment Canada, Ottawa, Ontario, 17 pp. + maps.

SEDGWICK, J.K., 1966. Geomorphology and mass budget of Peyto Glacier, Alberta. M.A. thesis, Department of Geography, McMaster University, Hamilton, Ontario, 165 pp.

SEDGWICK, J.K., 1967a. Peyto Glacier: a compendium of historical information. Glaciology Subdivision, Inland Waters Branch, Department of Energy, Mines and Resources, Ottawa, Ontario, 22 pp.

SEDGWICK, J.K., 1967b. Peyto Glacier: a compendium of recessional information. Unpublished report, Glaciology Subdivision, Inland Waters Branch, Department of Energy, Mines and Resources, Ottawa, Ontario, 29 pp.

SEDGWICK, J.K. and HENOCH, W.E.S., 1970. Peyto Glacier map. 1st Edition, Scale 1:10,000, Inland Waters Branch, Department of Energy, Mines and Resources, Ottawa, Ontario, map

SEDGWICK, J.K. and HENOCH, W.E.S., 1975. Peyto Glacier: general information. 2nd Edition, Scale 1:10,000, Glaciology Division, Water Resources Branch, Inland Waters Directorate, Environment Canada, Ottawa, Ontario, booklet, 30 pp. + map

SEDGWICK, J.K. and HENOCH, W.E.S., 1976. Peyto Glacier. Glaciology Division, Water Resources Branch, Inland Waters Directorate, Environment Canada, Ottawa, Ontario, map and booklet, 30 pp.

SMITH, N.D., 1978. Sedimentation processes and patterns in a glacier-fed lake with low sediment input. Canadian Journal of Earth Sciences, 15(5), p 741-756.

SMITH, N.D., 1981. The effect of changing sediment supply on sedimentation in a glacier-fed lake. Arctic and Alpine Research, 13(1), p 75-82.

SMITH, N.D. and SYVITSKI, J.P.M., 1982. Sedimentation in a glacier-fed lake: the role of pelletization on deposition of fine-grained suspensates. Journal of Sedimentary Petrology, 52(2), June, p 503-514.

SMITH, N.D., VENDL, M.A. and KENNEDY, S.K., 1982. Comparison of sedimentation regimes in four glacier-fed lakes of western Alberta. In - Research in Glacial, Glacio-fluvial and Glacio-lacustrine systems, Proceedings of 6th Guelph Symposium on Geomorphology, 9-10 May 1980, R. Davidson-Arnott, W. Nickling and B.D. Fahey, (Editors), Geographical Publication No.6, Department of Geography, University of Guelph, Geo Books Norwich, Geo Abstracts, Norwich, p 203-238.

STANLEY, A.D., 1970. A pilot study for an inventory of the glaciers in the Rocky Mountains: Inventory of Glaciers in the Waputik Mountains. In: Perennial Ice and Snow Masses, UNESCO-IASH Technical Papers in Hydrology No. 1. p 36-46.

STANLEY, A.D. and BELLAAR-SPRUYT, T., 1969. Combined mass, water and energy balance studies at Peyto Glacier. In - Proceedings of the North Saskatchewan Headwaters meeting, 5 December 1969, Ottawa, Inland Waters Branch, Department of Energy, Mines and Resources, Ottawa, Ontario, mimeo, unpaginated, 2 pp.

STENNING, A.J., 1980. Synoptic controls on katabatic layer characteristics above a temperate alpine-type glacier - Peyto Glacier, Alberta. M.Sc. thesis, Department of Geography, Memorial University of Newfoundland, St. John's, Newfoundland, August, 193 pp.

WALLACE, A.L., 1995. The Volumetric Change of the Peyto Glacier 1896-1966. MA. thesis, Department of Geography, Wilfrid Laurier University, Waterloo, Ontario, 141 pp.

YOUNG, G.J., 1970. Mass balance measurements related to surface geometry on Peyto Glacier, Alberta. In - Glaciers, Proceedings of IHD Workshop Seminar 1970, Vancouver, B.C., Canadian National Committee for the International Hydrological Decade, p 11-20.

YOUNG, G.J., 1974. Glacier mass-balance and terrain shape: a methodological study. Ph.D. thesis, Department of Geography, McGill University, Montreal, Quebec, 320 pp.

YOUNG, G.J., 1975. Accumulation and ablation patterns as functions of the surface geometry of a glacier. IAHS Publication No.104, p 134-138.

YOUNG, G.J., 1976. Discharge characteristics of a glacier-fed stream. Proceedings of CAGONT '76, Canadian Association of Geographers, Ontario Division Annual Meeting, 24 January 1976, Occasional Paper No.5, Department of Geography, Trent University, Peterborough, Ontario, p 42-43.

YOUNG, G.J., 1977. Relations between mass-balance and meteorological variables on Peyto Glacier, Alberta, 1967-1974. Zeitschrift fur Gletscherkunde und Glazialgeologie, 13(1/2), 111-125.

YOUNG, G.J., 1978. Influence of glaciers on streamflow. Plate 33, Hydrological Atlas of Canada, Fisheries and Environment Canada, Ottawa, Ontario, text and map.

YOUNG, G.J., 1981. The mass balance of Peyto Glacier, Alberta, Canada, 1965 to 1978. Arctic and Alpine Research, 13(3), August, p 307-318.

YOUNG, G.J., 1982. Hydrological relationships in a glacierized mountain basin. In - Hydrological Aspects of Alpine and High Mountain Areas, IAHS Publication No.138, p 51-59.

YOUNG, G.J., 1991. Hydrological interactions in the Mistaya basin, Alberta, Canada. In - Snow, Hydrology and Forests in High Alpine Areas, IAHS publication No.205, 237-244.

YOUNG, G.J., 1993. Melt Processes, terrain types and runoff generation in the Mistaya Basin, Alberta, Canada. Program and Abstracts, Scientific Meeting of the Canadian Geophysical Union, 9-11 May 1993, Banff, Alberta, Abstract No.15, unpaginated.

YOUNG, G.J. and ARNOLD, K.C., 1977a. Orthophotomaps of glaciers: an evaluation of an automated method applied to Peyto Glacier, Alberta. Zeitschrift fur Gletscherkunde und Glazialgeologie, 13(1/2), 99-110. Also - EΘS, Transactions, American Geophysical Union, 58(9), September, Abstract, p 902.

YOUNG, G.J. and ARNOLD, K.C., 1977b. A Stereo-orthophotograph of Peyto Glacier on the Gestalt Photomapper II. In - Fluctuations of glaciers 1970-1975 (Vol.III): a contribution to the International Hydrological Programme, Compiled for the Permanent Service on the Fluctuations of Glaciers of the IUGG-FAGS/ICSU by F. Muller, International Commission of Snow and Ice of the International Association of Hydrological Sciences, UNESCO, p 260-261.

YOUNG, G.J. and OMMANNEY, C.S.L., 1984. Canadian glacier hydrology and mass balance studies: a history of accomplishments and recommendations for future work. Geografiska Annaler, 66A(3), p 169-182.

YOUNG, G.J. and STANLEY, A.D., 1976. Canadian glaciers in the International Hydrological Decade program, 1965-1974. No.4: Peyto Glacier, Alberta - summary of measurements. Inland Waters Directorate Scientific Series No.71, Water Resources Branch, Fisheries and Environment Canada, Ottawa, Ontario, 59 pp.

Bow River

ALBERTA BOW RIVER WATER QUALITY TASK FORCE, 1991. Bow River: preserving our lifetime: report of the Bow River water quality task force. 78 pp.

ALBERTA BOW RIVER WATER QUALITY TASK FORCE, 1991. Bow River: preserving our lifetime: report of the Bow River water quality task force - Appendix vi. 116 pp.

ALBERTA ENVIRONMENT CONSERVATION AUTHORITY, 1974. Land use and resource development in the Eastern Slopes. Interim report #3: Bow River Basin. Edmonton Alberta. Environment Conservation Authority. Vol 7. 69 pp.

CHALLIES, J.B. and DENIS, L.G., 1916. Water Powers of Manitoba, Saskatchewan and Alberta/ Committee on Water and Water Powers, Commission of Conservation, Canada. Vol.9. 334 pp.

HAMILTON, H.R. and NORTH, L.J., 1986. Bow river water quality monitoring. Environmental protection services, pollution control division, Water quality control branch. 55 pp.

RUTTER, N.W., 1966. Glacial history of the Bow River Valley, Banff area, Alberta, Canadian Alpine Journal. Vol 49. p 157-173.

ULLMAN, P., 1972. Bow river water quality 1971-1972. Department of the Environment, Pollution Control Division. 25 pp.

WILSON, M., 1983. Once upon a river: archaeology and geology of the Bow River at Calgary, Alberta, Canada. Archaeological Survey of Canada. Ottawa, Natural Museums of Canada, Vol 35. 464 pp.

Historical Papers

BURPEE, L.J., 1914. Among the Canadian Alps. John Lane Company. New York. 239 pp.

FRASER, E., 1969. The Canadian Rockies Early Travels and Explorations. M.G.Hurtig LTD. Edmonton. 252 pp.

GROVE, J.M., 1988. The Little Ice Age. Metheun & Co. London. 498 pp.

HABEL, J., 1898. The North Fork of the Wapta. Appalachia, The Journal of the Appalachian Mountain Club, Vol 8. p 327-337.

LANG, A.H., 1943. Glaciers of the Rockies and Selkirks. Canadian Geographical Journal. Vol XXVI, No 2. p 56-67.

OUTRAM, J., 1905. In The Heart of The Canadian Rockies. Macmillan Company. New York. 464 pp.

SPRY, I.M., 1963. The Palliser Expedition: An Account of John Palliser's British North American Expedition 1857-1860. Macmillan company of Canada. Toronto. 620 pp.

SPRY, I.M., 1968. The Papers of the Palliser Expedition 1857-1860. The Champlain Society. Toronto. 680 pp.

STUTFIELD, H.E.M., and COLLIE, J.N., 1903. Climbs and exploration in the Canadian Rockies. Longmans,Green & Company. New York. 330 pp.

THOMPSON, C.S., 1898. At the Headwaters of the Bow. Appalachia. 8: p 320-327.

THORINGTON, J.M., 1925. The Glittering Mountains of Canada - Old Trails of the Rockies. John W.Lea Publishing. Philadelphia. 302 pp.

VAUX.Jr, G., and VAUX, W.S., 1907. Glacier Observations. Canadian Alpine Journal. Vol.1 No.1, Calgary, Alberta. p 138-148.

WALLACE, J.N., 1961. Early Exploration along the Bow and Saskatchewan Rivers. Alberta Historical Review. Spring, p 12-21.

WHEELER, A.O., 1933. Records of glacial observations in the Canadian Cordillera, 1933 and 1934. Canadian Alpine Journal, 22, p 172-187.

WILCOX, W.D., 1916. The Rockies of Canada. Knickerbocker Press. New York. 300 pp.

Appendix **Specific yield from sub-basins June, July, August, September, 1973-76**


Pipestone
(BA002)
Hector Lake
(BA008)
Baker Creek
(BA007)
Lake Louise
(BA001)
Johnson Creek
(BA006)
Forty Mile Creek
(BB003)
Standardized Discharge
(m³/sec/km²)
No Data
0.001-0.024
0.025-0.049
0.050-0.074
0.075-0.099
0.100-0.125
Banff
Redearth Creek
(BB005)
N
0
10
km
Brewster Creek
(BB004)

A.1 June 1973


Pipestone
(BA002)
Hector Lake
(BA008)
Baker Creek
(BA007)
Lake Louise
(BA001)
Johnson Creek
(BA006)
Forty Mile Creek
(BB003)
Standardized Discharge
(m³/sec/km²)
No Data
0.001-0.024
0.025-0.049
0.050-0.074
0.075-0.099
0.100-0.125
Banff
Redearth Creek
(BB005)
N
0
10
km
Brewster Creek
(BB004)

A.2 June 1974


Pipestone
(BA002)
Hector Lake
(BA008)
Baker Creek
(BA007)
Lake Louise
(BA001)
Johnson Creek
(BA006)
Forty Mile Creek
(BB003)
Standardized Discharge
(m³/sec/km²)
No Data
0.001-0.024
0.025-0.049
0.050-0.074
0.075-0.099
0.100-0.125
Banff
Redearth Creek
(BB005)
N
0
10
km
Brewster Creek
(BB004)

A.3 June 1975


Pipestone
(BA002)
Baker Creek
(BA007)
Hector Lake
(BA008)
Lake Louise
(BA001)
Johnson Creek
(BA006)
Forty Mile Creek
(BB003)
Standardized Discharge
(m³/sec/km²)
No Data
0.001-0.024
0.025-0.049
0.050-0.074
0.075-0.099
0.100-0.125
Banff
Redearth Creek
(BB005)
N
0
10
km
Brewster Creek
(BB004)

A.4 June 1976


Pipestone
(BA002)
Hector Lake
(BA008)
Baker Creek
(BA007)
Lake Louise
(BA001)
Johnson Creek
(BA006)
Forty Mile Creek
(BB003)
Standardized Discharge
(m³/sec/km²)
No Data
0.001-0.024
0.025-0.049
0.050-0.074
0.075-0.099
0.100-0.125
Banff
Redearth Creek
(BB005)
N
0
10
km
Brewster Creek
(BB004)

A.5 July 1973


Pipestone
(BA002)
Hector Lake
(BA008)
Baker Creek
(BA007)
Lake Louise
(BA001)
Johnson Creek
(BA006)
Forty Mile Creek
(BB003)
Standardized Discharge
(m³/sec/km²)
No Data
0.001-0.024
0.025-0.049
0.050-0.074
0.075-0.099
0.100-0.125
Banff
Redearth Creek
(BB005)
N
0
10
km
Brewster Creek
(BB004)

A.6 July 1974


Pipestone
(BA002)
Baker Creek
(BA007)
Hector Lake
(BA008)
Lake Louise
(BA001)
Johnson Creek
(BA006)
Forty Mile Creek
(BB003)
Standardized Discharge
(m³/sec/km²)
No Data
0.001-0.024
0.025-0.049
0.050-0.074
0.075-0.099
0.100-0.125
Banff
Redearth Creek
(BB005)
N
0
10
km
Brewster Creek
(BB004)

A.7 July 1975


Pipestone
(BA002)
Hector Lake
(BA008)
Baker Creek
(BA007)
Lake Louise
(BA001)
Johnson Creek
(BA006)
Forty Mile Creek
(BB003)
Standardized Discharge
(m³/sec/km²)
No Data
0.001-0.024
0.025-0.049
0.050-0.074
0.075-0.099
0.100-0.125
Banff
Redearth Creek
(BB005)
N
0
10
km
Brewster Creek
(BB004)

A.8 July 1976



A.9 August 1973



A.10 August 1974



A.11 August 1975



A.12 August 1976



A.13 September 1973



A.14 September 1974



A.15 September 1975



A.16 September 1976